

July 1983

RKER.83.120

GEOCHEMICAL INVESTIGATION OF A GAS SAMPLE FROM
WELL NOOKS FARM-1A, UNITED KINGDOM

by

J.M.A. Buiskool Toxopeus and J. Posthuma

Investigation 95.4600

This Report has been prepared by Shell in connection with its own internal evaluations and decision-making processes. Shell, its affiliates and each of their respective officers, employees and representatives make no representation or warranty whatsoever as to the accuracy or completeness of the information contained herein, and expressly disclaim any and all liability based on or relating to any information contained in, or errors or omissions from, this Report or based on or relating to the use of this Report. Shell will not enter into any correspondence relating to the contents of this Report.

KONINKLIJKE/SHELL EXPLORATIE EN PRODUKTIE LABORATORIUM

RIJSWIJK, THE NETHERLANDS

(Shell Research B.V.)

CONTENTS

	<u>Page</u>
Results and Conclusions	1

GEOCHEMICAL INVESTIGATION OF A GAS SAMPLE FROM
WELL NOOKS FARM-1A, UNITED KINGDOM

1. Results and Conclusions

A geochemical investigation has been carried out on a gas sample from well Nooks Farm-1A, United Kingdom. The composition of the gas is as follows

(mol %):

C ₁	82.60
C ₂	3.49
C ₃	0.94
iC ₄	0.10
nC ₄	0.19
iC ₅	0.04
nC ₅	0.04
C ₆ ⁺	0.06
CO ₂	0.19
N ₂	12.40
H ₂ s	-
CO	-

This composition and the carbon isotope value of -45.0 ‰ indicate that the gas is a gas of thermal origin generated from a kerogenous source rock, rather than a bacterial gas or gas from a humic source.

INITIAL DISTRIBUTION

3 Copies

Shell Expro London UEE/35/3/12

3 Copies

SIPM-EP 11/13